

The induced Chern-Simons term at finite temperature

L.L. Salcedo*

Departamento de Física Moderna, Universidad de Granada, E-18071 Granada, Spain

(Dated: February 1, 2008)

It is argued that the derivative expansion is a suitable method to deal with finite temperature field theory, if it is restricted to spatial derivatives only. Using this method, a simple and direct calculation is presented for the radiatively induced Chern-Simons-like piece of the effective action of (2+1)-dimensional fermions at finite temperature coupled to external gauge fields. The gauge fields are not assumed to be subjected to special constraints, and in particular, they are not required to be stationary nor Abelian.

PACS numbers: 11.10.Wx, 11.10.Kk, 11.15.-q, 11.30.Er

Keywords: Finite temperature; Chern-Simons; Three dimensions; Effective action; Gauge invariance

I. INTRODUCTION

Finite temperature field theory [1] is notoriously more difficult to deal with than its zero temperature counterpart. Lorentz symmetry is reduced and time and space play different roles. This is apparent in the imaginary time formalism, in which the time is Wick-rotated and compactified to a circle. As a consequence, tools such as the perturbative expansion or the derivative expansion, that were quite useful at zero temperature may become unreliable. These remarks apply, in particular, to (2 + 1)-dimensional QED. In this theory, due to the compactification, topologically non-trivial (large) gauge transformations are supported at finite temperature [2]. In principle, one would expect that the coefficient of the radiatively induced Chern-Simons (CS) term in the effective action [3] would be correctly quantized [4], consistently with gauge invariance of the partition function at any temperature [5]. This turns out not to be the case when that coefficient is computed using perturbation theory [2, 6]. Existing (Abelian and non-Abelian) exact results for particular configurations [5, 7, 8, 9] show that gauge invariance is an exact symmetry but perturbation theory fails to see this; perturbation theory is designed to describe the effective action in the neighborhood of a vanishing gauge field and large gauge transformations necessarily move the gauge field configuration away from the perturbative region. The derivative expansion is also problematic as already shown in a (0 + 1)-dimensional setting [10] (using the real time formalism). The problem is clear in the imaginary time formalism since the energy becomes a discrete variable. This suggests that the trouble exists only if one insists in taking a derivative expansion in both time and space. Technically, to make a derivative expansion of the effective action functional W , means to replace the given configuration $A_\mu(x)$ by a family of configurations $A_\mu(\lambda x)$, where λ is a book-keeping parameter, and then expand $W(\lambda)$ in powers of λ [21]. At finite temperature, this procedure is inconsis-

tent, since the external (bosonic) fields are required to be periodic in the (Euclidean) time and the dilatation in the temporal direction breaks this constraint. On the other hand, no problem should appear if the derivative expansion is restricted to the spatial directions (or at least, no new pathologies as compared to the zero temperature case). Moreover, gauge invariance can be accounted for if the bookkeeping parameter is introduced in the form $A_0(x_0, \lambda \mathbf{x})$, $\lambda A_i(x_0, \lambda \mathbf{x})$.

The exact results in [5, 8, 9] were obtained for particular configurations of the gauge field using specially adapted methods. In this Letter we show that general configurations are also amenable to explicit calculation using the method of expanding the effective action in the number of spatial covariant derivatives. We present a direct calculation of the induced CS term at finite temperature which yields a simple explicit expression for this quantity. This expression holds for general configurations and agrees with all previously known exact results. Of course, in addition to this, the effective action contains further contributions which are completely regular (i.e. gauge and parity invariant and ultraviolet finite) which are also calculable within the same general scheme.

The fermionic effective action is defined as $W = -\log \text{Det} D$, where $D = \gamma_\mu D_\mu + m$ is the Dirac operator, m is the fermion mass and $D_\mu = \partial_\mu + A_\mu$ is the covariant derivative [22]. The Dirac operator acts on a single-particle Hilbert space which contains space-time, Dirac and internal degrees of freedom (i.e. those associated to the gauge group). The finite temperature $T = 1/\beta$ is introduced by compactifying the Euclidean time. As usual the fermionic wave functions are antiperiodic and the gauge fields are periodic in the time variable.

As is well known, the effective action is afflicted by several related pathologies, namely, ultraviolet divergences, many-valuation and anomalies in some of the classical symmetries of the Dirac operator. Our purpose is to isolate precisely those pieces of the effective action which can have an anomalous or many-valued contribution. This excludes ultraviolet finite pieces, which are always regular. The expansion in the number of spatial covariant derivatives [11, 12] is appropriate for this kind of calculations, since this expansion preserves gauge in-

*Electronic address: salcedo@ugr.es

variance order by order and, in addition, terms beyond second order are ultraviolet finite. A further simplification can be achieved by selecting only those terms which have abnormal parity, i.e., those containing a Levi-Civita pseudo-tensor, since the normal parity component of the effective action can be renormalized preserving all classical symmetries and is one-valued [23]. In passing we note that “parity” (defined as space reflection but including $m \rightarrow -m$) is a symmetry of the classical action. Therefore, we would expect the normal (abnormal) parity component of the effective action to be an even (odd) function of the mass. This symmetry may be anomalously broken as a consequence of gauge invariance, as in the massless case [3].

II. THE CURRENT

Following Schwinger [13], we will work with the current, which is better behaved than the effective action. The current is defined through the relation $\delta W = \int d^3x \text{tr}(J_\mu \delta A_\mu)$ (where δA_μ is a local variation, $J_\mu(x)$ is a matrix in internal space and tr refers to this space), and thus

$$J_\mu(x) = -\text{tr}_D \langle x | \gamma_\mu \frac{1}{\gamma_\nu D_\nu + m} | x \rangle. \quad (1)$$

(tr_D refers to Dirac space.) In view of our previous remarks, our approach will be to compute the relevant terms of the spatial current J_i and then reconstruct the effective action from it. This requires to expand the current in powers of D_i retaining only terms with ϵ_{ij} and with precisely one D_i . A suitable way to deal with the matrix element at coincident points, which combines well with gradient expansion, is to use the method of symbols [14], adapted to finite temperature [11, 12] and improved by Pletnev and Banin [15, 16]. This gives

$$J_i(x) = -\frac{1}{\beta} \sum_{p_0} \int \frac{d^2 \mathbf{p}}{(2\pi)^2} \text{tr}_D \langle x | \gamma_i \frac{1}{\gamma_\nu \tilde{D}_\nu + m} | 0 \rangle. \quad (2)$$

In this formula $|0\rangle$ is the constant wave function, i.e. $\langle x|0\rangle = 1$ (which is periodic rather than antiperiodic), $p_0 = 2\pi(n + \frac{1}{2})/\beta$ runs over the fermionic Matsubara frequencies, and finally, $\tilde{D}_\mu$ is related to the original covariant derivative through a double similarity transformation,

$$\tilde{D}_\alpha = e^{i\partial_i^p D_i} e^{-ix_\mu p_\mu} D_\alpha e^{ix_\nu p_\nu} e^{-i\partial_j^p D_j}, \quad (\partial_i^p = \frac{\partial}{\partial p_i}). \quad (3)$$

The inner similarity transformation yields $D_\mu \rightarrow D_\mu + ip_\mu$ and corresponds to the original method of symbols. In that method, the integration over $\mathbf{p}$ cancels all contributions except those where D_i appears inside a commutator (more precisely, in the form $[D_i, \cdot]$). In the improved method, this cancellation is achieved prior to momentum

integration by means of the outer similarity transformation in (3). Explicitly:

$$\begin{aligned} \tilde{D}_0 &= ip_0 + D_0 - iE_i \partial_i^p + \frac{1}{2} [D_i, E_j] \partial_i^p \partial_j^p + O(D_i^3), \\ \tilde{D}_i &= ip_i + \frac{i}{2} F_{ij} \partial_j^p + O(D_i^3), \end{aligned} \quad (4)$$

where $F_{\mu\nu} = [D_\mu, D_\nu]$ and $E_i = F_{0i}$ is the electric field. The improvement does not extend to the compactified time coordinate, and indeed at finite temperature D_0 appears in two different ways which preserve gauge invariance, i) inside commutators, $[D_0, \cdot]$, and ii) through the (untraced) Polyakov loop $\Omega(x) = T \exp \left(- \int_{x_0}^{x_0+\beta} A_0(t, \mathbf{x}) dt \right)$ [12].

For our present purposes it will be sufficient to retain in (4) only those terms with at most one spatial covariant derivative. Upon substitution of these expressions in (2), and taking the Dirac trace keeping only the abnormal parity terms, produces

$$J_i^-(x) = 2im\epsilon_{ij} \frac{1}{\beta} \sum_{p_0} \int \frac{d^2 \mathbf{p}}{(2\pi)^2} \langle x | \frac{1}{\Delta} E_j \frac{1}{\Delta} | 0 \rangle + O(D_i^3), \quad (5)$$

where $\Delta = m^2 + \mathbf{p}^2 + (p_0 - iD_0)^2$. It is noteworthy that the result is directly ultraviolet finite and also consistent, i.e. a true variation [24].

The current (5) picks up contributions from all terms with two spatial indices in the effective action, however, most of those terms are perfectly regular, in the sense that they are ultraviolet finite, one-valued, strictly gauge invariant and anomaly free, therefore we can simplify the calculation by isolating only the anomalous terms, namely

$$J_i^{\text{an}} = im\epsilon_{ij} \frac{1}{\beta} \sum_{p_0} \int \frac{d^2 \mathbf{p}}{(2\pi)^2} \langle x | \left\{ \frac{1}{\Delta^2}, E_j \right\} | 0 \rangle. \quad (6)$$

This “anomalous” current is still consistent. The difference between the two currents, $J_i^- - J_i^{\text{an}}$, involves $[D_0, E_i]$ and derives from (fully calculable) regular terms in the effective action which are quadratic in E_i [17]. On the contrary the anomalous current does not contain such a commutator and so it cannot derive from one of those terms. After carrying out the momentum integration and the sum over frequencies, the anomalous current takes the simple explicit form

$$J_i^{\text{an}} = \epsilon_{ij} \langle x | \{ \varphi(D_0), E_j \} | 0 \rangle, \quad (7)$$

where we have introduced the function

$$\varphi(a) = \frac{i}{8\pi} \frac{\sinh(\beta m)}{\cosh(\beta m) + \cosh(\beta a)}. \quad (8)$$

This current preserves parity (as defined above), and can be written in a manifestly gauge invariant form by using the identity $e^{-\beta D_0} = \Omega(x)$ [12]:

$$J_i^{\text{an}} = \frac{i}{4\pi} \epsilon_{ij} \left\{ \frac{\sinh(\beta m)}{2 \cosh(\beta m) + \Omega + \Omega^{-1}}, E_j \right\}. \quad (9)$$

Note that this current gives the static limit of the gauge field self-energy instead of the long-wave limit [18]. The latter limit follows from restoring the regular terms stripped from the anomalous piece. Our criterion for selecting the anomalous piece is that the remainder is strictly gauge invariant, however, regarding the self-energy after rotation to real time, the anomalous piece is analytical at zero momentum and energy and the non-analytical behavior is contained in the remainder [17].

At this step we can already verify that the zero temperature limit comes out correctly. Indeed, for large β

$$J_i^{\text{an}} = \frac{i}{4\pi} \epsilon(m) \epsilon_{ij} E_j, \quad (T = 0) \quad (10)$$

(where $\epsilon(m)$ stands for the sign of m). This current derives from

$$W_{\text{an}} = -\frac{1}{2} \epsilon(m) W_{\text{CS}}, \quad (T = 0), \quad (11)$$

where W_{CS} is the Chern-Simons action

$$W_{\text{CS}} = \frac{i}{4\pi} \int d^3x \epsilon_{ij} \text{tr}(A_0 F_{ij} - A_i \partial_0 A_j). \quad (12)$$

Eq. (11) is the standard induced Chern-Simons term at zero temperature [2, 6, 19].

III. THE EFFECTIVE ACTION

Using the techniques developed in [11, 12] it is possible to reconstruct the anomalous piece of the effective action from its current while preserving manifest gauge invariance at every step of the calculation. This method will be presented elsewhere [17]. A more direct approach, which we will follow here, is to proceed by fixing the gauge [25]. We choose the gauge so that $A_0(x)$ is stationary

$$\partial_0 A_0(x) = 0. \quad (13)$$

Such a gauge always exists (globally) [11]. In this gauge, (9) becomes

$$J_i^{\text{an}}(x) = \epsilon_{ij} \{ \varphi(A_0(\mathbf{x})), E_j(x) \}. \quad (14)$$

The final step is to recover the effective action associated to this current. To this end we split the current into two parts $J_i^{(1)} + J_i^{(2)}$, corresponding to the decomposition of the electric field into $E_i = [D_0, A_i] - \partial_i A_0$. Noting that in our gauge A_0 and D_0 commute, it is readily verified that the first piece derives from

$$W^{(1)} = \int d^3x \epsilon_{ij} \text{tr}(\varphi(A_0) A_i [D_0, A_j]). \quad (15)$$

Thus the remainder must satisfy

$$\delta W^{(2)} = - \int d^3x \epsilon_{ij} \text{tr}(\delta A_i \{ \varphi(A_0), \partial_j A_0 \}) \quad (16)$$

To proceed, we further restrict the gauge (by means of a suitable subsequent stationary gauge transformation) so that $A_0(\mathbf{x})$ is everywhere a diagonal matrix, i.e

$$[A_0(\mathbf{x}), \partial_i A_0(\mathbf{x})] = 0. \quad (17)$$

For subsequent reference we note that in this gauge

$$[D_0, [D_0, A_i]] = [D_0, E_i]. \quad (18)$$

With this choice of gauge, (16) has the simple solution

$$W^{(2)} = - \int d^3x \epsilon_{ij} \text{tr}(\Phi(A_0) \partial_i A_j) \quad (19)$$

provided that $\Phi'(a) = 2\varphi(a)$ or, in our case,

$$\Phi(a) = \frac{i}{2\pi\beta} \tanh^{-1}(\tanh(\beta m/2) \tanh(\beta a/2)). \quad (20)$$

Using the convenient notation of differential forms (for the spatial indices only), the final result obtained by adding up (15) and (19), takes the form

$$W_{\text{an}} = \int dx_0 \text{tr}(\Phi(A_0)(A^2 - B) + \varphi(A_0) A[D_0, A]) \quad (21)$$

(where $A = A_i dx_i$ and $B = dA + A^2 = \frac{1}{2} F_{ij} dx_i dx_j$ is the magnetic field). This action reproduces the spatial components of the current. Since any further contributions to the effective action cannot contain A_i , they would be of the form $\int dx_0 \text{tr}(f(A_0) dA_0)$ and this vanishes identically in our gauge.

A subtle, but important, point is that, depending on the space manifold and the gauge group, a given configuration $A_0(\mathbf{x})$ may not be globally diagonalizable (i.e. with continuity at all points). Nevertheless, the “diagonal” gauge (17) always exists locally, that is, for each of the patches covering the space manifold. Without loss of generality we can assume that the support of the local variation in (16) takes place inside one of the patches, thus integration by parts in $\mathbf{x}$ is allowed and the correct current is obtained from (19). Because the current (14) is patch independent, this suggests that the integrand in (21) is also patch independent. As shown in the next section, this is actually correct, confirming that the functional W_{an} is well defined. For simplicity, we will momentarily assume $A_0(\mathbf{x})$ to be diagonal in a global gauge.

Eqs. (10) and (11) show that the calculation is consistent with the known zero temperature limit, and this can also be verified directly from W_{an} . Next we can study the behavior of this functional under gauge transformations. We have to distinguish between the “gauge fixing” transformation needed to bring the original configuration to the gauge where A_0 is stationary and diagonal, on the one hand, and the allowed gauge transformations which preserve the gauge conditions (13) and (17), on the other. Of course, gauge invariance of the partition function under the gauge fixing transformation cannot

be studied using our functional. Such gauge invariance follows from general arguments, e.g. by using the ζ -function version of the effective action [5]. The gauge conditions (13) and (17) are preserved by two kinds of gauge transformations, namely, stationary and discrete transformations [11]. The first class is that of transformations which are stationary and diagonal, and it is easily verified that W_{an} is invariant under such transformations. The second class refers to those transformations of the form $U = \exp(x_0 \Lambda)$, where Λ is a constant diagonal matrix with entries $\lambda_j = 2\pi i n_j / \beta$ (with n_j integer). Under discrete transformations A_i transforms covariantly (i.e., homogeneously) whereas $A_0 \rightarrow A_0 + \Lambda$. The current (14) transforms covariantly (due to periodicity of $\varphi(a)$) thus W_{an} can only change by constant additive terms, that is, terms that do not give a contribution to the transformed current. This can be verified directly from the functional as follows. First, at a formal level the function $\Phi(a)$ is periodic and W_{an} is invariant. More correctly, on its Riemann surface $\Phi(A_0 + \Lambda) = \Phi(A_0) + i\epsilon(m)\Lambda/4\pi$ and this induces a variation $-i\epsilon(m)/4\pi \int dx_0 \text{tr}(\Lambda dA)$ on W_{an} . This variation is precisely the same as that computed for the zero temperature functional in (11) under the same gauge transformation. This implies that this variation is of the form $\pm i\pi k n$, where k is an integer depending on the gauge group and the gauge field configuration and n is the winding number of the gauge transformation [26]. It can be noted that a perturbative expansion in W_{an} or J_i^{an} would break periodicity of the functions φ and Φ and, in general, this spoils gauge invariance under discrete transformations.

If there are large transformations and k is odd, the partition function changes by a factor $(-1)^n$ and gauge invariance must be enforced by adding a polynomial term $\pm \frac{1}{2} W_{\text{CS}}$ [3]. Such term is present directly in the effective action when the ζ -function regularization is used [11, 19]. This term is even as a function of m and so it introduces an anomalous breaking of parity. Note that this new term also adds a parity breaking contribution to the current; although the current (7) is gauge and parity invariant it does not derive from a gauge invariant partition function. It also noteworthy that for massless fermions the factor $(-1)^n$ is also present [3] but comes from the normal parity component of the effective action, since W_{an} vanishes in this case.

In [5, 8] it is shown that the full (abnormal parity) result is $-\int dx_0 \text{tr}(\Phi(A_0)B)$ provided that the configuration is stationary, Abelian and A_0 is $\mathbf{x}$ -independent. The same formula is found in [9] in the non-Abelian stationary case when the electric field vanishes. This is consistent with our W_{an} noting that in both cases the A_i are stationary and commute everywhere with A_0 , and so the explicit A cancel in (21).

IV. TOPOLOGICAL ISSUES

Let us now analyze the general case in which the diagonal gauge is not assumed to exist globally. In this case it will be convenient to work in the global gauge where A_0 is stationary but not necessarily diagonal, and rewrite W_{an} in that gauge. This yields

$$W_{\text{an}} = \int dx_0 \text{tr}(\Phi(A_0)(\mathcal{A}^2 - B) + \varphi(A_0) \mathcal{A}[D_0, \mathcal{A}]) \quad (22)$$

In this formula $\mathcal{A}$ denotes the vector field A of the diagonal gauge rotated covariantly back to the stationary gauge. That is, if U denotes the stationary gauge transformation which brings A_0 to a (local) diagonal gauge,

$$\mathcal{A} = A + dUU^{-1}. \quad (23)$$

A crucial property of (22) is that it remains unchanged under the replacement

$$\mathcal{A} \rightarrow \mathcal{A}' = \mathcal{A} + C, \quad (24)$$

provided that the field $C(x)$ satisfies

$$\partial_0 C = [A_0, C] = 0. \quad (25)$$

This is readily verified noting that C is a 1-form and A_0 , D_0 and C are all commuting quantities.

By construction $\mathcal{A}$ transforms covariantly (under transformations of the global stationary gauge and fixed local diagonal gauge), but is not globally defined since the gauge transformation U only exists locally. Let $M^{(k)}$ denote a set of local charts covering the space manifold. In each chart we can take a diagonalizing gauge transformation $U^{(k)}$ and this defines a corresponding ‘‘covariant vector field’’ $\mathcal{A}^{(k)}$. What have to be verified is that 2-form in (22) takes the same value in any of the charts.

To show this let us note that the relation (18), which was written in the diagonal gauge, in the stationary gauge becomes

$$[D_0, [D_0, \mathcal{A}^{(k)}]] = [D_0, E]. \quad (26)$$

Therefore the field $C^{(k,\ell)} = \mathcal{A}^{(k)} - \mathcal{A}^{(\ell)}$ (which is stationary due to (23)) satisfies

$$[A_0, [A_0, C^{(k,\ell)}]] = 0. \quad (27)$$

This already implies that $C^{(k,\ell)}$ commutes with A_0 and satisfies (25). As a consequence any two $\mathcal{A}^{(k)}$ at the same point are related as in (24) and W_{an} is patch independent.

It is noteworthy that the field $A'_0(\mathbf{x})$, denoting A_0 in a diagonal gauge, is (or can be taken to be) equal in all patches. Indeed, the local eigenvalues of $A_0(\mathbf{x})$, $a_\lambda(\mathbf{x})$, can be labeled at each point so that they are continuous functions on the space manifold. Thus, without loss of generality we can choose the local diagonal gauges in such a way that the eigenvalues are ordered in all patches in the same way and so the $A'_0(\mathbf{x})$, which are diagonal

matrices containing the eigenvalues, will be the same matrix in all patches. However, although $A'_0(\mathbf{x})$ is global, it needs not correspond to a global gauge; in general, the eigenvectors corresponding to $a_\lambda(\mathbf{x})$ cannot be chosen in a continuous way on the space manifold. Assuming the non degenerated case for simplicity, the choice of eigenvectors in each patch will differ by a phase (and perhaps normalization, depending on the gauge group). These phases are contained in the diagonal transition matrices $U^{(\ell)-1}U^{(k)}$, which contain the non trivial topology of $A_0(\mathbf{x})$ (under gauge transformations).

When we bring a completely general gauge field configuration to an A_0 -stationary gauge the result is not unique. The different possible stationary gauges so obtained are related either by a stationary gauge transformation or by a discrete transformation [11]. The discrete transformations are of the form $U = \exp(x_0\Lambda)$, where $\Lambda(\mathbf{x})$ is stationary, commutes with $A_0(\mathbf{x})$ and has eigenvalues $\lambda_j = 2\pi i n_j/\beta$ (with n_j integer). The discussion of how W_{an} changes under these transformations is essentially the same as that given before for the case of a global diagonal gauge. It can be noted, however, that the discrete transformations need not be topologically small if the diagonal gauge is not global (an explicit $SU(2)$ ex-

ample is shown in [2], also discussed in [11].)

V. CONCLUSIONS

In summary, we have shown that the method of expanding in the number of spatial covariant derivatives is suitable to make computations at finite temperature, and in particular it is able to retain all subtleties usually tied to the effective action, such as topological pieces, many-valuation and anomalies. At the practical level, the method have been shown to be quite efficient, yielding a simple explicit form for the induced CS term at finite temperature, without assuming special constraints on the gauge field configuration.

Acknowledgments

This work is supported in part by funds provided by the Spanish DGICYT grant no. PB98-1367 and Junta de Andalucía grant no. FQM-225.

-
- [1] N. P. Landsman and C. G. van Weert, Phys. Rept. **145**, 141 (1987).
 - [2] R. D. Pisarski, Phys. Rev. **D35**, 664 (1987).
 - [3] A. N. Redlich, Phys. Rev. Lett. **52**, 18 (1984).
 - [4] S. Deser, R. Jackiw, and S. Templeton, Ann. Phys. **140**, 372 (1982).
 - [5] S. Deser, L. Griguolo, and D. Seminara, Phys. Rev. Lett. **79**, 1976 (1997), hep-th/9705052.
 - [6] K. S. Babu, A. K. Das, and P. Panigrahi, Phys. Rev. **D36**, 3725 (1987).
 - [7] G. Dunne, K.-Y. Lee, and C.-h. Lu, Phys. Rev. Lett. **78**, 3434 (1997), hep-th/9612194.
 - [8] C. Fosco, G. L. Rossini, and F. A. Schaposnik, Phys. Rev. Lett. **79**, 1980 (1997), hep-th/9705124.
 - [9] F. T. Brandt, A. K. Das, and J. Frenkel (2001), hep-th/0107120.
 - [10] J. Barcelos-Neto and A. K. Das, Phys. Rev. **D58**, 085022 (1998), hep-th/9805204.
 - [11] L. L. Salcedo, Nucl. Phys. **B549**, 98 (1999), arXiv:hep-th/9802071.
 - [12] C. Garcia-Recio and L. L. Salcedo, Phys. Rev. **D63**, 045016 (2001), arXiv:hep-th/0007183.
 - [13] J. S. Schwinger, Phys. Rev. **82**, 664 (1951).
 - [14] L. L. Salcedo and E. Ruiz Arriola, Ann. Phys. **250**, 1 (1996), arXiv:hep-th/9412140.
 - [15] N. G. Pletnev and A. T. Banin, Phys. Rev. **D60**, 105017 (1999), arXiv:hep-th/9811031.
 - [16] L. L. Salcedo, Eur. Phys. J. **C20**, 161 (2001), arXiv:hep-th/0012174.
 - [17] L. L. Salcedo (2002), in preparation.
 - [18] Y.-C. Kao and M.-F. Yang, Phys. Rev. **D47**, 730 (1993).
 - [19] R. E. Gamboa Saravi, G. L. Rossini, and F. A. Schaposnik, Int. J. Mod. Phys. **A11**, 2643 (1996), hep-th/9411238.
 - [20] A. N. Sisakian, O. Y. Shevchenko, and S. B. Solganik, Nucl. Phys. **B518**, 455 (1998), hep-th/9710022.
 - [21] This shows that each term of the expansion is independent of the *method* used in the calculation. This fails to hold for other expansions, such as the commutator expansion.
 - [22] Our conventions follow those in [11, 12] with $\eta = +1$.
 - [23] This remark applies to the massive case only. In the massless case [3], or if $|\mu| > |m|$ (where μ is the chemical potential) [20], the normal parity term combines with the abnormal parity one to yield a one-valued partition functional DetD .
 - [24] Consistency requires $\int d^3x \text{tr}(\delta_1 A_i \delta_2 J_i)$ to be symmetric under the exchange of the labels 1 and 2.
 - [25] The gauge conditions (13) and (17) depend only on A_0 and so they do not interfere with an arbitrary local variation of $A_i(x)$, as required to obtain the current and reconstruct the effective action from it.
 - [26] As introduced here, discrete transformations are always topologically small for simply connected groups, however, this is not necessarily the case when the diagonal gauge is not global (see section IV).